

SMALL MODELS OF GRAPH COLOURING MANIFOLDS AND THE STIEFEL MANIFOLDS $\text{Hom}(C_5, K_n)$

CARSTEN SCHULTZ

ABSTRACT. We show Péter Csorba's conjecture that the graph homomorphism complex $\text{Hom}(C_5, K_{n+2})$ is homeomorphic to a Stiefel manifold, the space of unit tangent vectors to the n -dimensional sphere. For this a general tool is developed that allows to replace the complexes $\text{Hom}(G, K_n)$ by smaller complexes that are homeomorphic to them whenever G is a graph for which those complexes are manifolds. The equivariant version of Csorba's conjecture is proved up to homotopy.

We also study certain subdivisions of simplicial manifolds that are related to the interval poset of their face posets and their connection with geometric approximations to diagonal maps.

1. INTRODUCTION

The study of properties of graphs, especially their chromatic number, through topological spaces associated to the graphs began with Lovász' proof of Kneser's conjecture [Lov78]. Recently the focus has shifted to the homomorphism complex $\text{Hom}(G, H)$ associated to two graphs G and H , see [BK06a]. It is a cell complex whose vertices are the graph homomorphisms from G to H and whose topology captures the way in which homomorphisms can be transformed into each other by local changes; its cells correspond to multi-homomorphisms from G to H , functions which assign to every vertex of G a set of vertices of H such that every choice function for it is a homomorphism, see Definition 2.4. In particular Babson and Kozlov have proved a conjecture of Lovász that states that if for a graph G and an $r \geq 1$ the complex $\text{Hom}(C_{2r+1}, G)$ is $(n-1)$ -connected then G is not $(n+2)$ -colourable [BK06b, Sch05]. The proof uses the functoriality of Hom and topological properties of the complexes $\text{Hom}(C_{2r+1}, K_{n+2})$.

Despite these advances, the homotopy or homeomorphism types of very few of the complexes $\text{Hom}(G, H)$ are known, even in the case where H is a complete graph and graph homomorphisms become colourings. The cohomology groups of the spaces $\text{Hom}(C_m, K_{n+2})$ have been calculated in [Koz05a]. Susequent to this and the current work it has been established in [Sch06] that the colimit of a diagram of all $\text{Hom}(C_m, K_{n+2})$ with even m is homotopy equivalent to the free loop space of $\mathbb{S}^n$ while the colimit over all odd m is homotopy equivalent to the space of all loops in $\mathbb{S}^n$ which are equivariant with respect to the antipodal actions on $\mathbb{S}^1$ and $\mathbb{S}^n$.

Among the spaces $\text{Hom}(C_m, K_{n+2})$, the spaces $\text{Hom}(C_5, K_{n+2})$ are special, because they are manifolds. Spaces $\text{Hom}(G, K_n)$ which are manifolds, graph colouring manifolds, have been studied by Csorba and Lutz [CL05]. Based on cohomology and

Date: October 2005, introduction updated in July 2006.

This research was supported by the Deutsche Forschungsgemeinschaft within the European graduate program "Combinatorics, Geometry, and Computation" (No. GRK 588/2).

index calculations, Péter Csorba has conjectured that $\text{Hom}(C_5, K_{n+2})$ is homeomorphic to the Stiefel manifold of orthonormal 2-frames in $\mathbb{R}^{n+1}$ [Cso05, Conj. 4.8]. A proof of this is the main result of this article.

Theorem (5.2). *Let $n \geq 0$. Then there is a homeomorphism*

$$(1) \quad \text{Hom}(C_5, K_{n+2}) \approx \{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle = 0\}.$$

The case $n = 0$, $\text{Hom}(C_5, K_2) = \emptyset$, is trivial, since C_5 admits no 2-colouring. The case $n = 1$, $\text{Hom}(C_5, K_3) \approx \mathbb{S}^1 \times \mathbb{S}^0$, is easily checked. The cases $n = 2$ and $n = 3$ are proved in [CL05], $\text{Hom}(C_5, K_4) \approx \mathbb{R}P^3$ by direct construction, and $\text{Hom}(C_5, K_5) \approx \mathbb{S}^3 \times \mathbb{S}^2$ by a one week computer calculation.

Our proof consists of two distinct parts, and this article is structured accordingly. The first part consists of finding a smaller model for $\text{Hom}(C_5, K_{n+2})$ and is presented in Section 3. If I is a set of vertices of a graph G , then one can define a complex $\text{Hom}_I(G, H)$ that is derived from $\text{Hom}(G, H)$ by identifying graph homomorphisms that differ only on vertices in I . If I is an independent set of vertices, then $\text{Hom}_I(G, H)$ is homotopy equivalent to $\text{Hom}(G, H)$. This has first been proved and used in [Cso05, Sec. 2.8]. We show that for graph colouring manifolds it does indeed yield homeomorphic complexes.

Lemma (3.9, 3.12). *Let G be a graph and I an independent set of vertices of G . If $\text{Hom}(G, K_n)$ is a manifold for all n , then $\text{Hom}(G, K_n) \approx \text{Hom}_I(G, K_n)$ for all n .*

We hope that this result will find further applications. At least the fact alone that $\text{Hom}_I(G, H)$ is a complex with a smaller number of vertices than $\text{Hom}(G, H)$ should make the determination of the homeomorphism types of more graph colouring manifolds accessible to computer calculations just Péter Csorba has used the homotopy equivalence of $\text{Hom}(G, H)$ and $\text{Hom}_I(G, H)$ to, among other things, simplify computer calculations of cohomology groups of homomorphism complexes. In the current work, however, this construction is used to expose the structure of $\text{Hom}(C_5, K_{n+2})$.

Let I be a maximal independent set of vertices of C_5 . Removing it from C_5 leaves us with an edge and an isolated vertex. The colourings of the edge give rise to an n -sphere, since $\text{Hom}(K_2, K_{n+2}) \approx \mathbb{S}^n$. Similarly the colourings of the isolated vertex together with the information that this vertex is not isolated in C_5 give rise to another n -sphere. It follows that $\text{Hom}_I(C_5, K_{n+2})$ can be identified with a subspace of $\mathbb{S}^n \times \mathbb{S}^n$. It is not the full space $\mathbb{S}^n \times \mathbb{S}^n$, since not every multi-colouring of the edge and the isolated vertex can be extended to C_5 ; the edge and the isolated vertex cannot be regarded separately but interact via the vertices in I . It turns out that $\text{Hom}_I(C_5, K_{n+2})$ is the common boundary of regular neighbourhoods of two subspaces of $\mathbb{S}^n \times \mathbb{S}^n$, each corresponding to an element of I . These subspaces are ambiently isotopic to the subspaces $\{(x, x) : x \in \mathbb{S}^n\}$ and $\{(x, -x) : x \in \mathbb{S}^n\}$, which determines $\text{Hom}_I(C_5, K_{n+2})$ up to homeomorphism and proves the Theorem. The relevant subdivisions of the spheres and constructions of regular neighbourhoods are examples of general constructions that can be carried out for any pl-triangulation of a manifold, the triangulation being the boundary of an $(n+1)$ -simplex in the case of $\text{Hom}_I(C_5, K_{n+2})$. These constructions are carried out in Section 4.

Section 5 puts together the results from the two preceding sections. In it we also consider Csorba's stronger conjecture, also stated in [Cso05], that the homeomorphism of (1) can be chosen in such a way that it transports the involution on $\text{Hom}(C_5, K_{n+2})$ that appears in the Theorem of Babson and Kozlov mentioned above to the involution $(x, y) \mapsto (x, -y)$ on $\mathbb{S}^n \times \mathbb{S}^n$. Our current approach yields

only a proof of a homotopy version of this conjecture, the complete conjecture remains open.

Acknowledgements. I would like to thank Péter Csorba for comments on a draft of this work. These helped to clarify some points and lead to the inclusion of the part on the involution on $\text{Hom}(C_5, K_{n+2})$.

2. PRELIMINARIES

We collect some definitions, notations and results that we will need.

Posets and cellular complexes. For a partially ordered set, or *poset*, P , we denote by ΔP its *order complex*, the simplicial complex with vertex set P that consists of all chains in P . A monotone (or antitone) map $f: P \rightarrow Q$ between posets induces a simplicial map $\Delta f: \Delta P \rightarrow \Delta Q$. All cell complexes are assumed to be regular. For a cell complex C we denote its *face poset* by FC and its underlying space by $|C|$. Thus $\Delta(FC)$ is the barycentric subdivision of the complex C and $|\Delta(FC)| \approx |C|$. If C and D are cell complexes, then $C \times D$ denotes the cell complex whose face poset is isomorphic to $FC \times FD$, even if C and D are simplicial complexes. Since most of our simplicial complexes are order complexes, we have no need for a notation for the simplicial complex that is their product, $\Delta(P \times Q)$ is the usual simplicial subdivision of the cell complex $\Delta P \times \Delta Q$. Our preferred way to construct homotopies between maps between order complexes is the following special case of [Seg68, Prop. 2.1], which we will use without citing it.

2.1. Lemma. *Let P, Q be posets. If $f, g: P \rightarrow Q$ are order preserving functions such that $f(p) \leq g(p)$ for all $p \in P$, then the maps $\Delta f, \Delta g: \Delta P \rightarrow \Delta Q$ are homotopic.*

Proof. The map $H: \{0, 1\} \times P \rightarrow Q$ defined by $H(0, x) := f(x)$, $H(1, x) := g(x)$ is order preserving and hence yields the desired homotopy

$$I \times \Delta P \approx \Delta(\{0, 1\} \times P) \xrightarrow{\Delta H} \Delta Q,$$

where we view $\{0, 1\}$ as a poset. □

For a poset P its *dual poset* P^o is the poset with the same elements but the order reversed. When considering $p \in P$ as an element of P^o we write it as p^o . Thus $p \mapsto p^o$ is an antitone bijection $P \rightarrow P^o$.

For a cell c , its dimension is denoted by $|c|$, the cardinality of a set M is written as $\#M$. When identifying a simplex σ with the set of its vertices we thus have $|\sigma| = \#\sigma - 1$.

Graphs and graph complexes. All Graphs that we consider are finite, simple, and without loops. The vertex set of a graph G is denoted by $V(G)$, and for $S \subset V(G)$ the set of all common neighbours of the elements of S is denoted by $\nu(S)$. A graph homomorphism from G to H is a function $f: V(G) \rightarrow V(H)$ that respects the edge relation, i.e. such that $f[\nu(\{v\})] \subset \nu(\{f(v)\})$ for all $v \in V(G)$.

2.2. Definition. For a graph G we denote by $\text{ind}(G)$ the poset of all independent subsets of $V(G)$, including the empty set, ordered by inclusion. By $\text{Ind}(G)$ we denote the *independence complex* of G , i.e. the simplicial complex with vertex set $V(G)$ and simplices the independent subsets of $V(G)$.

2.3. Definition. Let G, H be graphs. A *multi-homomorphism* from G to H is a function $\varphi: V(G) \rightarrow \mathcal{P}(V(H)) \setminus \{\emptyset\}$ such that every function $f: V(G) \rightarrow V(H)$ with $f(v) \in \varphi(v)$ for all $v \in V(G)$ is a graph homomorphism.

2.4. Definition. Let G, H be graphs. A function $\varphi: V(G) \rightarrow \mathcal{P}(V(H)) \setminus \{\emptyset\}$ can be identified with a cell of the cell complex

$$\prod_{v \in V(G)} \Delta^{\#V(H)-1}.$$

The subcomplex of all cells indexed by multi-homomorphisms is denoted by $\text{Hom}(G, H)$. We identify elements of $F\text{Hom}(G, H)$ with the corresponding multi-homomorphisms. If $f: G' \rightarrow G, g: H \rightarrow H'$ are graph homomorphisms, then there is an induced monotone map

$$\begin{aligned} F\text{Hom}(f, g): F\text{Hom}(G, H) &\rightarrow F\text{Hom}(G', H'), \\ (F\text{Hom}(f, g))(\varphi)(v) &:= g[\varphi(f(v))] \end{aligned}$$

and hence a continuous map

$$|\text{Hom}(f, g)| \rightarrow |\text{Hom}(G, H)| \rightarrow |\text{Hom}(G', H')|.$$

The above constructions are functorial. Details can be found in [BK06a].

2.5. Notation. Let $n \in \mathbb{N}$. P_n denotes that path with n edges on the vertex set $\{0, \dots, n\}$. K_n denotes the complete graph on n vertices with vertex set $\{1, \dots, n\}$. C_n is the cycle of length n with vertex set $\{1, \dots, n\}$.

Piecewise linear topology.

2.6. Definition. Let Y be a simplicial complex and X a subcomplex of Y . The subcomplex

$$N := \{\sigma \in Y: \text{There is a } \tau \in X \text{ with } \sigma \cup \tau \in Y\}$$

of Y is called the *simplicial neighbourhood of X in Y* and

$$\dot{N} := \{\sigma \in N: \sigma \cap \tau = \emptyset \text{ for all } \tau \in X\}$$

is called the *simplicial neighbourhood boundary*. We also call the pair $(N, \dot{N})$ the simplicial neighbourhood.

We will need the following form of the Simplicial Neighbourhood Theorem [RS82, 3.11].

2.7. Theorem. *Let M be a pl-triangulated compact manifold, X a full subcomplex of M , and $(N, \dot{N})$ the simplicial neighbourhood of X in M . If $(N, \dot{N})$ is a manifold with boundary, then N is a regular neighbourhood of X in M .*

3. RESTRICTION MAPS OF Hom-COMPLEXES

Given graphs G and H and an independent subset I of $V(G)$ there is a subcomplex of $\text{Hom}(G \setminus I, H)$ which is homotopy equivalent to $\text{Hom}(G, H)$. The use of this smaller complex in the study of the homotopy type of $\text{Hom}(G, H)$ has been introduced in [Cso05, Sec. 2.8]. The same idea has been implicitly used in [Sch05] to study the complexes $\text{Hom}(C_{2r+1}, K_{n+2})$.

In Lemma 3.5 we give a criterion that allows us to detect cases in which the two spaces are actually homeomorphic. We will use this to study $\text{Hom}(C_5, K_{n+2})$.

We round off the investigation by showing that the applicability of the Lemma to C_5 is not incidental but a consequence of the fact that $\text{Hom}(C_5, K_n)$ is a manifold.

For the results on $\text{Hom}(C_5, K_{n+2})$, only Definition 3.1, Lemma 3.5, and Example 3.8 will be needed from this section.

Homotopy equivalences. We recall the definition of $\text{Hom}_S(G, H)$ from [Cso05]. The proofs and examples given here are for expository purposes only. Neither are they needed for the homeomorphism results we present afterwards, nor do we claim any originality here, except perhaps for the point of view described in Example 3.4.

3.1. Proposition and Definition ([Cso05]). *Let G, H be graphs and $S \subsetneq V(G)$. Let $i: G \setminus S \rightarrow G$ denote the inclusion of the subgraph induced on the complement of S . We consider the continuous map*

$$\text{Hom}(i, H): \text{Hom}(G, H) \rightarrow \text{Hom}(G \setminus S, H)$$

and denote its image by $\text{Hom}_S(G, H)$.

- (i) *The map $\text{Hom}(i, H)$ is cellular and $\text{Hom}_S(G, H)$ is a subcomplex of the cell-complex $\text{Hom}(G \setminus S, H)$.*
- (ii) *If S is independent, then $\text{Hom}_S(G, H)$ consists of all cells of $\text{Hom}(G \setminus S, H)$ indexed by multihomomorphisms φ with $\nu\left(\bigcup_{u \in \nu(\{v\})} \varphi(u)\right) \neq \emptyset$ for all $v \in S$.*
- (iii) *If S is independent, then the map*

$$\text{Hom}(G, H) \rightarrow \text{Hom}_S(G, H)$$

induced by $\text{Hom}(i, H)$ is a homotopy equivalence.

Proof. The first two claims are obvious. Let us denote the inclusion $S \rightarrow G$ by j . Then

$$(F\text{Hom}(i, H), F\text{Hom}(j, H)): F\text{Hom}(G, H) \rightarrow F\text{Hom}(G \setminus S, H) \times F\text{Hom}(S, H)$$

is an embedding. If S is independent, then the image of this embedding is

$$\{(\varphi, \rho) \in \text{im } F\text{Hom}(i, H) \times F\text{Hom}(S, H) : \rho \leq a(\varphi)\}$$

where a is the antitone map $\text{im } F\text{Hom}(i, H) \rightarrow F\text{Hom}(S, H)$ given by

$$a(\varphi)(v) := \nu\left(\bigcup_{u \in \nu(\{v\})} \varphi(u)\right).$$

The result follows from Lemma 4.3 □

We illustrate the Proposition by deriving two well-known facts from it.

3.2. Example. Let H be a graph, T be a tree with at least three vertices and v a leaf of T . Since any multihomomorphism from $T \setminus \{v\}$ can be extended to one from T by mapping v to one of the vertices assigned to one of the neighbours in $T \setminus \{v\}$ of the neighbour of v , the restriction map $\text{Hom}(T, H) \rightarrow \text{Hom}(T \setminus \{v\}, H)$ is surjective and hence a homotopy equivalence. By induction any inclusion $i: K_2 \rightarrow T$ of an edge in a tree induces a homotopy equivalence $\text{Hom}(T, H) \simeq \text{Hom}(K_2, H)$. This example generalizes to folds in the first parameter of Hom , see [Koz05b] for this concept.

3.3. Example. $\text{Hom}(K_2, H)$ is one of the constructions called the *box complex* of H , while $\text{Hom}(K_1, H)$ is just the full simplex on the vertices of H . Considering an inclusion $i: K_1 \rightarrow K_2$, the image of the map $\text{Hom}(i, H)$ consists of the simplices corresponding to sets $A \subset V(H)$ with $\nu(A) \neq \emptyset$. This is the *neighbourhood complex* of H introduced in [Lov78]. Thus the box complex is homotopy equivalent to the neighbourhood complex. This example also appears in [Cso05]. For comparison of several graph complexes also see [MZ, CLSW04, Cso04, Živ05].




FIGURE 1.






FIGURE 2.

We look more closely at $\text{Hom}(P_2, K_{n+2})$. This will give us an opportunity to introduce some ideas that can be generalised, but are not needed for the remainder of this article.

3.4. Example. The relevant inclusions related to maximal independent subsets of P_2 are $j: K_1 \cong \{1\} \rightarrow P_2$ and $k: P_2 \setminus \{1\} \rightarrow P_2$. For $M \subset V(K_{n+2})$, $M \neq \emptyset$, we define $X_M := \{\varphi \in F\text{Hom}(P_2, K_{n+2}): \varphi(1) = M\}$ and $Y_M := \{\varphi \circ k: \varphi \in X_M\}$. The sets $\Delta(Y_{\{u\}})$, $u \in V(K_{n+2})$ cover $\text{im Hom}(k, K_{n+2})$, and $Y_M \cap Y_N = Y_{M \cup N}$ for all M, N . Also $Y_{V(K_{n+2})} = \emptyset$. So, if we define the poset $Q := \mathcal{P}(V(K_{n+2})) \setminus \{\emptyset, V(K_{n+2})\}$ ordered by reverse inclusion, then Y is a diagram of spaces over Q , where the maps are given by inclusion, and $\text{Hom}_{\{1\}}(P_2, K_{n+2}) \approx \text{colim } Y$. For the example $n = 1$ the poset Q and the diagram Y are shown in Figure 1 and Figure 2, and $\text{Hom}_{\{1\}}(P_2, K_3)$ is depicted in Figure 3. On the other hand $\text{Hom}(P_2, K_{n+2})$, which is shown in Figure 4, can be seen to be homeomorphic to $\text{hcolim } Y$, and $\text{Hom}(k, K_{n+2})$ corresponds to the natural map from $\text{hcolim } Y$ to $\text{colim } Y$ which is a homotopy equivalence by the Projection Lemma, since Y is a free diagram because of its construction from a cover. This indicates another way in which Proposition 3.1 could be proved. We also see that in general $\text{Hom}(G, H)$ will not be homeomorphic to $\text{Hom}_I(G, H)$, even if I is independent.

The diagram Y can also be used to describe the map $\text{Hom}(j, K_{n+2})$ as the map from the homotopy colimit of Y to the homotopy colimit of the constant diagram over Q which assigns a point to every element. The homotopy colimit of this constant diagram is just ΔQ , which is homeomorphic to $\mathbb{S}^n$. This map is a homotopy equivalence, because all Y_M are contractible, which can be seen to be a consequence of the independence of the set $\{0, 2\}$. This description is nearer to the proof of Proposition 3.1 given above. In general the poset Q would have to be replaced by the face poset of $\text{Hom}_S(G, H)$. The situation in this example is special, because P_2 is bipartite.

Homeomorphisms. We examine situations in which the complexes $\text{Hom}(G, K_n)$ and $\text{Hom}_I(G, K_n)$ are indeed homeomorphic. The independence complex of G (Definition 2.2) plays an important role in recognising these situations.

FIGURE 3. $\text{Hom}_{\{1\}}(P_2, K_3)$ FIGURE 4. $\text{Hom}(P_2, K_3)$

3.5. Lemma. *Let G be a graph, $n \in \mathbb{N}$, and $S \subsetneq V(G)$ an independent set. We define subsets A_v of $\text{ind}(G)$ for all $v \in V(G)$ by*

$$A_v := \{I \in \text{ind}(G) : v \in I\}$$

and similarly

$$B_v := \begin{cases} \{I \in \text{ind}(G \setminus S) : v \in I\}, & v \notin S \\ \{I \in \text{ind}(G \setminus S) : I \cup \{v\} \in \text{ind}(G)\}, & v \in S. \end{cases}$$


If there is a homeomorphism $h: |\Delta(\text{ind}(G))| \rightarrow |\Delta(\text{ind}(G \setminus S))|$ with $h[|\Delta(A_v)|] = |\Delta(B_v)|$ for all $v \in V(G)$, then $\text{Hom}(G, K_n) \approx \text{Hom}_S(G, K_n)$.

Proof. We look at $\text{Hom}(G, K_n)$ one colour at a time instead of one vertex at a time. The map

$$\begin{aligned} \gamma: F\text{Hom}(G, K_n) &\rightarrow \prod_{c \in V(K_n)} \text{ind}(G) \\ \varphi &\mapsto (\{u \in V(G) : c \in \varphi(u)\})_c \end{aligned}$$

is a well-defined poset embedding. Hence $\text{Hom}(G, K_n)$ is homeomorphic to the order complex of the image of γ . The poset $\text{ind}(G)^{V(K_n)}$ describes multicolourings without the condition that each vertex has to obtain at least one colour. Spelling out this additional condition yields

$$\text{im } \gamma = \bigcap_{v \in V(G)} \bigcup_{d \in V(K_n)} \prod_{c \in V(K_n)} \begin{cases} A_v, & c = d \\ \text{ind}(G), & c \neq d. \end{cases}$$

FIGURE 5. $\Delta(\text{ind } C_5)$ vs. $\Delta(\text{ind}(C_5 \setminus \{2, 4\}))$

Since the A_v are closed from above, the operation of taking the order complex commutes with all constructions appearing in this equation, and

$$\text{Hom}(G, K_n) \approx \bigcap_{v \in V(G)} \bigcup_{d \in V(K_n)} \prod_{c \in V(K_n)} \begin{cases} \Delta(A_v), & c = d \\ \Delta(\text{ind}(G)), & c \neq d. \end{cases}$$

follows. In the same way $F\text{Hom}(G \setminus S, K_n)$ embeds in $\text{ind}(G \setminus S)^{V(K_n)}$. $\text{Hom}_S(G, K_n)$ is distinguished in this embedding by the condition that for every vertex in S one of the colours is not used by any of its neighbours. Since B_v is closed from above for $v \in S$ and closed from below for $v \notin S$, the above argument can be repeated to obtain

$$\text{Hom}_S(G, K_n) \approx \bigcap_{v \in V(G)} \bigcup_{d \in V(K_n)} \prod_{c \in V(K_n)} \begin{cases} \Delta(B_v), & c = d \\ \Delta(\text{ind}(G \setminus S)), & c \neq d. \end{cases}$$

It is now obvious how the homeomorphism h induces a homeomorphism $\text{Hom}(G, K_n) \approx \text{Hom}_S(G, K_n)$. $\square$

Continuing Example 3.3 we give a quite trivial application of the Lemma.

3.6. Example. We consider the inclusion $i: K_1 \rightarrow K_2$. $\text{ind}(K_2) = \{\emptyset, \{0\}, \{1\}\}$, hence $\Delta(\text{ind}(K_2))$ is homeomorphic to an interval. The two points of the boundary are $\Delta(\{\{0\}\}) = \Delta(A_0)$ and $\Delta(\{\{1\}\}) = \Delta(A_1)$. On the other hand $\text{ind}(K_1) = \{\emptyset, \{0\}\}$, so $\Delta(\text{ind}(K_1))$ is also homeomorphic to an interval. The boundary points are $\Delta(\{\{0\}\}) = \Delta(B_0)$ and $\Delta(\{\{1\}\}) = \Delta(B_1)$. Thus the Lemma is applicable, and $\text{Hom}(K_2, K_{n+2}) \approx \text{Hom}_{\{1\}}(K_2, K_{n+2})$. The latter complex is the boundary of an $(n+1)$ -simplex. This gives yet another proof of $\text{Hom}(K_2, K_{n+2}) \approx \mathbb{S}^n$.

3.7. Example. The Lemma cannot be applied to the restriction maps considered in Example 3.4 as we have seen there. Indeed $\Delta(\text{ind}(P_2))$ is not homeomorphic to $\Delta(\text{ind}(K_1))$, $\Delta(\text{ind}(K_2))$, or $\Delta(\text{ind}(P_2 \setminus \{1\}))$. $\text{ind}(P_2)$ has the maximal elements $\{1\}$ and $\{0, 2\}$ and hence is not a pure complex. As we have seen $\Delta(\text{ind}(K_1))$ and $\Delta(\text{ind}(K_2))$ are intervals. $\text{ind}(P_2 \setminus \{1\}) \cong \text{ind}(K_1) \times \text{ind}(K_1)$ and hence $\Delta(\text{ind}(P_2 \setminus \{1\}))$ is homeomorphic to a 2-disk.

A less trivial example and our main reason for this investigation.

3.8. Example. We consider $\{2, 4\} \subset V(C_5)$. Figure 5 shows that Lemma 3.5 can be applied and hence $\text{Hom}(C_5, K_{n+2}) \approx \text{Hom}_{\{2, 4\}}(C_5, K_{n+2})$.

Manifolds. In the two examples we have just seen where $\text{Hom}(G, K_n) \approx \text{Hom}_I(G, K_n)$ for a non-empty independent set I and all n the independence complex $\text{Ind}(G)$ is a sphere, $\text{Ind}(K_2) \approx \mathbb{S}^0$ and $\text{Ind}(C_5) \approx \mathbb{S}^1$. This is what makes $\text{Hom}(G, K_n)$ a manifold in these cases. Complexes $\text{Hom}(G, K_n)$ that are manifolds have been investigated in [CL05]. We show that for these $\text{Hom}(G, K_n) \approx \text{Hom}_I(G, K_n)$ always holds.

3.9. Theorem ([CL05]). *Let G be a graph. The following statements are equivalent.*

- (i) $\text{Ind}(G)$ is a *pl-sphere*.
- (ii) $\text{Hom}(G, K_n)$ is a *pl-manifold* for all n .
- (iii) $\text{Hom}(G, K_{\chi(G)+1})$ is a *pl-manifold*.

If these statements hold, then $\dim \text{Hom}(G, K_n) = n(\dim \text{Ind}(G) + 1) - |V(G)|$.

We repeat the proof from [CL05], since we will afterwards build upon the ideas used in it.

Proof. Let $\varphi \in F \text{Hom}(G, K_n)$. For every $k \in V(K_n)$ the cell φ determines an independent set $\{v \in V(G) : k \in \varphi(v)\}$ and hence a simplex σ_k of $\text{Ind}(G)$, where we allow the empty simplex. The link of φ in $\text{Hom}(G, K_n)$ is homeomorphic to the join of the links of the σ_k in $\text{Ind}(G)$. If $\text{Ind}(G) \approx \mathbb{S}^m$, then the link of σ_k is an $(m - 1 - |\sigma_k|)$ -sphere, and the link of φ is a sphere of dimension

$$n - 1 + \sum_k (m - 1 - |\sigma_k|) = n(m + 1) - 1 - n - \sum_k |\sigma_k| = n(m + 1) - \#(V(G)) - 1 - |\varphi|,$$

so $\text{Hom}(G, K_n)$ is a manifold of dimension $n(m + 1) - \#(V(G))$. If $n > \chi(G)$ then there is a cell φ for which the empty simplex is among the σ_k , so the link of φ is a join of spaces of which one is $\text{Ind}(G)$. For the join to be a sphere it is necessary for $\text{Ind}(G)$ to be a sphere, see [RS82, 2.24(5)]. $\square$

We will want to apply Lemma 3.5 in this case and therefore start examining the relationship between $\text{Ind}(G)$ and $\text{Ind}(G \setminus S)$.

3.10. Lemma. *Let G be a graph and $S \subset V(G)$ a non-empty independent set. Then $\text{Ind}(G \setminus S)$ is the complement of the interior of the simplicial neighbourhood of the simplex S in $\text{Ind}(G)$. If $\text{Ind}(G)$ is a *pl-manifold*, then this simplicial neighbourhood is a regular neighbourhood. Hence if $\text{Ind}(G) \approx \mathbb{S}^n$ then $\text{Ind}(G \setminus S) \approx \mathbb{D}^n$.*

Proof. We only have to show that the simplicial neighbourhood of S is a regular neighbourhood if $\text{Ind}(G)$ is a manifold. We assume that $\text{Ind}(G)$ is an n -manifold. The simplicial neighbourhood of the simplex S consists of all $I \in \text{ind}(G) \setminus \{\emptyset\}$ such that there exists a $v \in S$ such that $I \cup \{v\} \in \text{Ind}(G)$. Its boundary consists of those I for which additionally $I \cap S = \emptyset$. By Theorem 2.7 it will be sufficient to show that the simplicial neighbourhood is a manifold with boundary, the boundary being as just described. To prove this we take a simplex I in the boundary and examine its link. Its link in $\text{Ind}(G)$ can be identified with $\text{Ind}(G')$ where G' is the graph obtained from G by deleting the vertices in I and all of their neighbours and is an $(n - \#I)$ -sphere, since $\text{Ind}(G)$ is an n -manifold. We set $S' := S \cap V(G')$. Since I is a simplex of the simplicial neighbourhood of S , S' is non-empty. In the link of I the boundary of the simplicial neighbourhood of S corresponds to the simplicial neighbourhood of S' in $\text{Ind}(G')$, which is an unlinked $(n - 1 - \#I)$ -sphere by induction. $\square$

3.11. Remark. Since $\text{Ind}(G)$ determines G , it is easy to describe, which simplicial complexes are independence complexes of graphs. These are the flag simplicial complexes. The preceding Lemma can therefore be viewed as a statement on flag complexes.

3.12. Lemma. *Let G be a graph, $n \in \mathbb{N}$, and $S \subsetneq V(G)$ an independent set. If $\text{Ind}(G)$ is a pl-sphere, then $\text{Hom}(G, K_n) \approx \text{Hom}_S(G, K_n)$.*

Proof. We want to apply Lemma 3.5 and use the notation introduced there. Let $m := \dim \text{Ind}(G)$. $\Delta(\text{ind}(G))$ is a cone over $\text{Ind}(G)$, therefore an $(m+1)$ -ball, and A_v is the star of v in the barycentric subdivision of $\text{Ind}(G)$, i.e. the cell of the dual complex which is dual to v . We will now describe $\Delta(\text{ind}(G \setminus S))$.

If $S \neq \emptyset$, then we have seen in the preceding Lemma that $\text{Ind}(G \setminus S)$ is an m -ball. This ball is covered by the sets $\Delta(B_v)$ with $v \in V(G) \setminus S$. The boundary of this ball consists of the simplices I for which there exists a $v \in S$ such that $I \cup \{v\}$ is independent. Therefore the sets $\Delta(B_v)$ with $v \in S$ cover the cone over this boundary in $\Delta(\text{ind}(G \setminus S))$. It follows that $\Delta(\text{ind}(G \setminus S))$ is an $(m+1)$ -ball and that the sets $\Delta(B_v)$ for all $v \in V(G)$ cover the boundary of this ball. The last statement is also true if $S = \emptyset$, and we will from here on allow this case in order to facilitate inductive arguments.

For $M \in \text{ind}(G) \setminus \{\emptyset\}$ we set

$$C_M := \{I \in \text{ind}(G \setminus S) : M \setminus S \subset I, I \cup M \in \text{ind}(G)\}.$$

Then $B_v = C_{\{v\}}$, and we have $\Delta(C_M) \cap \Delta(C_N) = \Delta(C_{M \cup N})$, if $M \cup N$ is independent, and $C_M \cap C_N = \emptyset$ otherwise.

Let $M \in \text{ind}(G) \setminus \{\emptyset\}$ and let G' be the graph obtained from G by removing all vertices in M and their neighbours. As seen in the proof of the preceding Lemma, $\text{Ind}(G')$ is homeomorphic to the link of M in $\text{Ind}(G)$ and hence an $(m - \#M)$ -sphere. Let $S' := S \cap V(G')$. The map

$$\begin{aligned} C_M &\rightarrow \text{ind}(G' \setminus S') \\ I &\mapsto I \setminus M \end{aligned}$$

is an isomorphism and for $N \supsetneq M$ it maps C_N to

$$\{I \in \text{ind}(G' \setminus S') : (N \setminus M) \setminus S \subset I, I \cup (N \setminus M) \in \text{ind}(G')\}.$$

Therefore by induction $\Delta(C_M)$ is an $(m - |M|)$ -ball and its boundary is the union of the $\Delta(C_N)$ for $N \supsetneq M$.

This shows that the sets $\Delta(C_M)$ form a cell-decomposition of the boundary of $\Delta(\text{ind}(G \setminus S))$ that has a face poset that is isomorphic to the face poset of the dual complex of the simplicial complex $\text{Ind}(G)$. This yields a homeomorphism from the boundary of the ball $\Delta(\text{ind}(G))$ to the boundary of the ball $\Delta(\text{ind}(G \setminus S))$ that carries $\Delta(A_v)$ to $\Delta(B_v)$ for every $v \in V(G)$. This homeomorphism can be extended to a homeomorphism between balls and Lemma 3.5 can be applied. $\square$

4. SUBDIVISIONS AND DIAGONAL APPROXIMATIONS

We examine several constructions that occur naturally in the study of homomorphism complexes of graphs, although they will not be mentioned in this section. All posets in this section are assumed to be finite.

Edge subdivision. For an ordered simplicial complex there is a subdivision that introduces a vertex for every edge, which is why we will call it its *edge subdivision*. For an order complex of a poset, it is the order complex of the interval poset. It has been first described in [Wal88]. In connection with graph complexes it has been used in [Živ05].

4.1. Definition and Proposition. *For a poset P we define the closed interval poset of P to be the subposet of $P \times P^\circ$ consisting of all (p, q°) with $p \leq q$ and denote it by $\text{Int } P$. Int is a functor from posets to posets. $\Delta(\text{Int } P)$ is a subdivision of ΔP . More precisely, choosing a point $x_{p,q}$ in the interior of the simplex $\langle p, q \rangle$ of ΔP for every pair (p, q) with $p < q$ determines a unique map $|\Delta(\text{Int } P)| \rightarrow |\Delta P|$ which sends (p, p°) to p , (p, q°) to $x_{p,q}$ for $p < q$, and is affine on simplices, and this map is a homeomorphism. In particular for $0 < \lambda < 1$ we define $h_P^\lambda: |\Delta \text{Int } P| \rightarrow |\Delta P|$ by choosing $x_{p,q}$ as $\lambda p + (1 - \lambda)q$. h^λ is a natural equivalence from $P \mapsto |\Delta \text{Int } P|$ to $P \mapsto |\Delta P|$. We abbreviate $h_P^{1/2}$ as h_P . All choices of $x_{p,q}$ yield isotopic maps, and any such map is homotopic to each of the two maps $\Delta(\text{Int } P) \rightarrow \Delta P$ given by $(p, q^\circ) \mapsto p$ or $(p, q^\circ) \mapsto q$.*

Proof. To see that $\Delta(\text{Int } P)$ is a subdivision of ΔP one either checks directly that the map $|\Delta(\text{Int } P)| \rightarrow |\Delta P|$ is bijective as in [Wal88], or one uses that for a simplex with vertices $p_0 < p_1 < \dots < p_r$ the poset $\text{Int } \{p_0, \dots, p_r\}$ has the minimum (p_0, p_r°) and the elements (p_0, q°) and (q', p_r°) are comparable only if one of them equals this minimum. Hence $\Delta(\text{Int } \{p_0, \dots, p_r\})$ is a cone over $\Delta(\text{Int } \{p_0, \dots, p_{r-1}\}) \cup \Delta(\text{Int } \{p_1, \dots, p_r\})$, which yields a recursive description of the subdivision. To connect subdivision maps for different choices of $x_{p,q}$ by an isotopy, one moves the points $x_{p,q}$. The homotopies are constructed by extending the definition of h_P^λ to $\lambda = 0$ and $\lambda = 1$. $\square$

4.2. Remark. $(p_0, q_0^\circ) \leq (p_1, q_1^\circ) \iff [p_0, q_0] \supset [p_1, q_1]$, hence the name interval poset.

For illustration we use this subdivision to prove a simple Lemma, which is quite specialised but useful in the proof of Proposition 3.1.

4.3. Lemma. *Let P, Q be posets, $a: P \rightarrow Q$ an antitone map, and*

$$R := \{(p, q) \in P \times Q : q \leq a(p)\}.$$

Assume that any two elements of Q having a common upper bound have a unique least upper bound. Then the map $\Delta(R) \rightarrow \Delta(P)$ induced by projection onto the first factor is a homotopy equivalence.

Proof. We define monotone maps $\pi: R \rightarrow P$, $(p, q) \mapsto p$ and $\sigma: \text{Int } P \rightarrow R$, $(p_0, p_1^\circ) \mapsto (p_0, a(p_1))$. Then $(\pi \circ \sigma)(p_0, p_1^\circ) = p_0$, so $|\pi| \circ |\sigma| \simeq h_P$. Also

$$(\sigma \circ \text{Int } \pi)((p_0, q_0), (p_1^\circ, q_1^\circ)) = (p_0, a(p_1)) \leq (p_0, a(p_1) \vee q_0) \geq (p_0, q_0)$$

and, since $((p_0, q_0), (p_1^\circ, q_1^\circ)) \mapsto (p_0, a(p_1) \vee q_0)$ is a well-defined monotone map, $|\sigma| \circ h_P^{-1} \circ |\pi| \circ h_R \simeq h_R$. Thus $|\sigma| \circ h_P^{-1}$ is homotopy inverse to $|\pi|$. $\square$

4.4. Remark. In the proof the chain poset of P would have worked equally well, but the interval poset seems more natural. Also we will see that iterated interval posets are easier to describe than iterated chain posets.

Diagonal approximation. By construction the edge subdivision of ΔP is a subcomplex of $\Delta(P \times P^o) \approx \Delta P \times \Delta P$. It can be viewed as a substitute for the diagonal, that is not a subcomplex of $\Delta(P \times P^o)$.

4.5. Proposition. *Let P be a poset and $o: \Delta P \rightarrow \Delta P^o$ denote the homeomorphism induced by $p \mapsto p^o$. Then the diagonal map $\Delta(\text{Int } P) \xrightarrow{h_P} \Delta P \xrightarrow{(\text{id}, \Delta o)} \Delta P \times \Delta P^o$ is isotopic to the map $\Delta(\text{Int } P) \rightarrow \Delta(P \times P^o) \approx \Delta P \times \Delta P^o$ induced by inclusion.*

Proof. For $0 \leq \lambda \leq 1$ we consider the map $(h_P^\lambda, \Delta o \circ h_P^{1-\lambda}): \Delta(\text{Int } P) \rightarrow \Delta P \times \Delta P^o$. For $\lambda = 1$ this map agrees with the map induced by inclusion and is hence an embedding. For $0 < \lambda < 1$ both components are homeomorphism, so it is also an embedding. For $\lambda = \frac{1}{2}$ the map agrees with the diagonal map. $\square$

Iterated and generalised edge subdivisions. Iterating the interval poset construction yields again easy to describe posets.

4.6. Proposition. *Let P be a poset. The map*

$$\begin{aligned} \text{Int}(\text{Int } P) &\rightarrow \{(p, q^o, r, s^o) \in P \times P^o \times P \times P^o : p \leq q \leq r \leq s\} \\ ((x, y^o), (u, v^o)^o) &\mapsto (x, u^o, v, y^o) \end{aligned}$$

is an isomorphism of posets. $\square$

This isomorphism makes it natural to expect that $\Delta\{(p, q^o, r) \in P \times P^o \times P : p \leq q \leq r\}$ would also be a subdivision of ΔP , and we will confirm this in the next Proposition. It also indicates another connection between $\Delta(\text{Int}(\text{Int } P))$ and $\Delta(\text{Int } P)$ than the first being the edge subdivision of the second. We therefore take a closer look at the complex $\Delta \text{Int } P$.

An r -simplex of $\Delta \text{Int } P$ is of the form $(p_0, q_0^o) < \dots < (p_r, q_r^o)$. This implies $p_0 \leq p_1 \leq \dots \leq p_r \leq q_r \leq q_{r-1} \leq \dots \leq q_0$. Hence the simplex is contained in $\Delta(\{p_i\} \times \{q_j^o\})$ and $\max\{p_i\} \leq \min\{q_j\}$. On the other hand for chains $p_0 < \dots < p_r$ and $q_0 < \dots < q_s$ with $p_r \leq q_0$ the subset $\{p_i\} \times \{q_i^o\}$ of $P \times P^o$ is contained in $\text{Int } P$. $\Delta(\{p_i\} \times \{q_i^o\})$ is the product of an r -simplex and an s -simplex and its image under $h_P: |\Delta \text{Int } P| \rightarrow |\Delta P|$ is contained in the simplex $\Delta(\{p_i\} \cup \{q_j\})$. Thus the sub-complex of the cell-complex $\Delta P \times \Delta P$ consisting of all cells $\langle p_0, \dots, p_r \rangle \times \langle q_0, \dots, q_s \rangle$ with $\max\{p_i\} \leq \min\{q_j\}$ is isomorphic to a cell-subdivision of ΔP , and it has $\Delta \text{Int } P$ as a simplicial subdivision. It follows that any subdivision of $\Delta P \times \Delta P$ leads to a subdivision of ΔP . In particular $\Delta \text{Int } P$ arises in this way by subdividing $\Delta P \times \Delta P$ as $\Delta(P \times P^o)$, and $\Delta(\text{Int}(\text{Int } P))$ arises by subdividing $\Delta P \times \Delta P$ as $\Delta(\text{Int } P \times \text{Int } P)$. On the other hand, the subdivision of ΔP obtained in this way by triangulating $\Delta P \times \Delta P$ as $\Delta(P \times P)$ is not a full subcomplex of $\Delta(P \times P)$, which underlines the usefulness of the interval poset.

4.7. Proposition. *Let P be a poset. There is a commutative diagram*

$$\begin{array}{ccc} |\Delta \text{Int } P| & \xrightarrow{\quad} & |\Delta(P \times P^o)| \\ \uparrow \approx & & \uparrow \approx \\ |\Delta\{(p, q^o, r) \in P \times P^o \times P : p \leq q \leq r\}| & \xrightarrow{\quad} & |\Delta(\text{Int } P \times P)| \end{array}$$

with the horizontal arrows induced by inclusion and the vertical arrows homeomorphisms.




FIGURE 6.

Proof. $\Delta(\text{Int } P \times P)$ is a subdivision of $\Delta P \times \Delta P$. According to the preceding discussion we only have to check that $\Delta\{(p, q^o, r) \in \text{Int } P \times P : q \leq r\}$ is the correct subcomplex. Let $(p_0, q_0^o, r_0) < \dots < (p_k, q_k^o, r_k)$ be a chain in $\text{Int } P \times P$. The smallest cell of $\Delta P \times \Delta P$ that contains it is $\Delta(\{p_i\} \cup \{q_i\}) \times \Delta\{r_i\}$. Now $\max(\{p_i\} \cup \{q_i\}) = q_0$, $\min\{r_i\} = r_0$, and $q_0 \leq r_0$ if and only if $q_i = r_i$ for all i . $\square$

4.8. Remark. While $\Delta \text{Int } P$ is a subdivision of ΔP and hence, as we have used, $\Delta(\text{Int } P \times P)$ and $\Delta(P \times P^o)$ are both subdivisions of $\Delta P \times \Delta P$, the complex $\Delta(\text{Int } P \times P)$ is not a subdivision of $\Delta(P \times P^o)$.


Manifolds. In general there is no ambient isotopy from the inclusion map $\Delta(\text{Int } P) \rightarrow \Delta(P \times P^o)$ to the diagonal map, as the case where ΔP is a simplex shows. If ΔP is a manifold however, then there is one.

4.9. Proposition. *Let P be a poset such that ΔP is a pl-triangulation of a compact manifold M^n . Then the isotopy from Proposition 4.5 is ambient.*

The technical part of the following proof is needed only for $n = 2$. In Theorem 5.2 this case is used for $\text{Hom}(C_5, K_4) \approx \mathbb{R}P^3$, which has already been proved in [CL05].

Proof. The codimension of M in $M \times M$ equals n . Since any isotopy in codimension at least 3 is ambient ([Zee63], see also [RS82, 7.3]) and a simple drawing (Figure 6) should convince the reader of the truth of the Proposition for $n = 1$, we will spell out the details only for $n = 2$.

Let $n = 2$. We will construct a sequence of subpolyhedra of $M \times M$, the first one being the diagonal, the last one being $\Delta(\text{Int } P)$, such that for any consecutive pair there is an isotopy of $M \times M$ which is equal to the identity outside a small neighbourhood of a 4-cell of the form $\sigma \times \sigma$ with σ a simplex of ΔP and carries one polyhedron into the other. Each of these subpolyhedra will be of the following form. For $p_0 < p_1 < p_2$, $p_i \in P$, it will contain either the diagonal of $\langle p_0, p_1, p_2 \rangle \times \langle p_0, p_1, p_2 \rangle$ or $\Delta(\text{Int } \{p_0, p_1, p_2\})$, i.e. the union of the simplices $\langle p_0 \rangle \times \langle p_0, p_1, p_2 \rangle$, $\langle p_0, p_1 \rangle \times \langle p_1, p_2 \rangle$, and $\langle p_0, p_1, p_2 \rangle \times \langle p_2 \rangle$. Additionally, for $p_0 < p_1$ such that for exactly one of the two 2-simplices adjacent to $\langle p_0, p_1 \rangle$ the polyhedron contains the corresponding part of the diagonal, the simplex $\langle (p_0, p_0), (p_0, p_1), (p_1, p_1) \rangle$ will be included. Now assume we are given two such subpolyhedra Q_0, Q_1 which differ only for the choice for the simplex $\langle p_0, p_1, p_2 \rangle$. We assume that Q_0 agrees on $D := \langle p_0, p_1, p_2 \rangle \times \langle p_0, p_1, p_2 \rangle$ with the diagonal, and that we know by induction

FIGURE 7. The cell-complex with face poset $\text{Int } M$.

that Q_0 is a locally flat submanifold of $M \times M$. Let N be a small closed regular neighbourhood of D . It suffices to show that both $(N, N \cap Q_i)$ are unknotted ball pairs, because there is then an isotopy of N which is constant on the boundary of N and moves $N \times Q_0$ to $N \times Q_1$ [RS82, 4.4, 3.22(i)], and that can be extended to $M \times M$ by the identity. Since Q_0 is a submanifold and we will see that $D \cap Q_0$ is a 2-ball with boundary $\overline{Q_0 \setminus D}$, we know that if N is chosen small enough, then $(N, N \cap Q_i)$ is homeomorphic to $(D, D \cap Q_i)$ with an external collar attached to $(\partial D, \overline{Q_0 \setminus D})$. Since Q_i is contained in the union of all cells of the form $\sigma \times \sigma$ with σ a simplex of ΔP and the intersection of two of these is either empty or also of this form, $\overline{Q_0 \setminus D}$ is the union of either $\langle p_i \rangle \times \langle p_i, p_j \rangle \cup \langle p_i, p_j \rangle \times \langle p_j \rangle$ or $\langle (p_i, p_i), (p_j, p_j) \rangle$ for every pair $i < j$. $D \cap Q_i$ is a two ball with boundary $\overline{Q_0 \setminus D}$. $D \cap Q_0$ consists of $\langle (p_0, p_0), (p_1, p_1), (p_2, p_2) \rangle$ plus simplices $\langle (p_i, p_i), (p_i, p_j), (p_j, p_j) \rangle$ as needed. $D \cap Q_1$ consists of $\Delta(\text{Int } \{p_0, p_1, p_2\})$ plus the remaining simplices of the form $\langle (p_i, p_i), (p_i, p_j), (p_j, p_j) \rangle$. It can therefore be checked without further knowledge about M that $(N, N \cap Q_i)$ is an unknotted $(4, 2)$ -ball pair for $i \in \{0, 1\}$. $\square$

If P is the face poset of a manifold M , then there is another useful way to see $\Delta(\text{Int } P)$, which is the edge subdivision of the barycentric subdivision of M , as a subdivision of a cell-complex. In this case P° is the face poset of the dual cell-complex M° , and hence $P \times P^\circ$ is the face poset of $M \times M^\circ$. $\Delta(\text{Int } P)$ is then the barycentric subdivision of a subcomplex of $M \times M^\circ$, namely the subcomplex of all cells $\sigma \times \tau^\circ$, where τ is a simplex of M , τ° its dual cell, and σ a face of τ . The maximal cells are those with $\sigma = \tau$. Thus $\Delta(\text{Int } P)$ is the barycentric subdivision of a cell-complex which is homeomorphic to M and contains a facet for every face of M . Figure 7 shows this cell-complex for a part of a manifold. Figure 6 shows it as a subcomplex of $M \times M^\circ$, where M is the boundary of a 2-simplex.


4.10. Proposition. *Let P be the face poset of a pl-triangulation of a compact manifold M^n . Let $N, B \subset P^\circ \times \text{Int } P$ be given by*

$$N := \{(p^\circ, q, r^\circ) : q \leq r, p \cap r \neq \emptyset\},$$

$$B := \{(p^\circ, q, r^\circ) : q \leq r, p \cap r \neq \emptyset, p \not\leq q\},$$

where notationally identified elements of P with sets of vertices. Then ΔN is a $2n$ -dimensional manifold with boundary ΔB .

Proof. Let $s = (p^\circ, q, r^\circ) \in P^\circ \times \text{Int } P$. Then $\Delta \{t \in P^\circ \times \text{Int } P : t < s\} \approx \mathbb{S}^{2n-|p|-|r|+q-1}$. So $P^\circ \times \text{Int } P$ is the face poset of a cell complex with N and B

FIGURE 8. The situation of Propositions 4.10 and 4.12 for $M = \mathfrak{d}\Delta^2$.

corresponding to subcomplexes, and it will be sufficient to describe the links of cells of this complex. We define

$$\begin{aligned} \text{lk}(s) &:= \{t \in P^o \times \text{Int } P : t > s\}, \\ \text{lk}^+(s) &:= \{(u^o, v, w^o) \in \text{lk}(s) : u \cap w \neq \emptyset\}, \\ \text{lk}^-(s) &:= \{(u^o, v, w^o) \in \text{lk}(s) : u \not\subset v\}. \end{aligned}$$

For $s \in N$ we will have to show that

$$(2) \quad (\Delta \text{lk}(s); \Delta \text{lk}^+(s), \text{lk}^-(s)) \approx \begin{cases} (\mathbb{S}^{d-1}; \mathbb{S}^{d-1}, \emptyset), & s \in N \setminus B \\ (\mathbb{S}^{d-1}; \mathbb{D}_+^{d-1}, \mathbb{D}_-^{d-1}), & s \in B \end{cases}$$

with $d = |r| - |q| + |p|$.

To be able to approach this inductively, we consider the link of s in a direction given by a set A of vertices. For $A \subset p \cup r$, we make the following auxiliary definitions.

$$\begin{aligned} \text{lk}_A(s) &:= \{(u^o, v, w^o) \in \hat{P}^o \times \text{Int } P : (u^o, v, w^o) > s, u \Delta p \subset A, v \Delta q \subset A, w \Delta r \subset A\}, \\ \text{lk}_A^+(s) &:= \{(u^o, v, w^o) \in \text{lk}_A(s) : u \cap w \cap A \neq \emptyset\}, \\ \text{lk}_A^-(s) &:= \{(u^o, v, w^o) \in \text{lk}_A(s) : u \cap A \not\subset v\}. \end{aligned}$$

Here $\hat{P}$ is P with a minimum, the empty simplex, added, and $u \Delta p$ denotes the symmetric difference $(u \cup p) \setminus (u \cap p)$.

For $A \cap B = \emptyset$ we have

$$\begin{aligned} (3) \quad & (\Delta \text{lk}_{A \cup B}(s); \Delta \text{lk}_{A \cup B}^+(s), \Delta \text{lk}_{A \cup B}^-(s)) \approx (\Delta \text{lk}_A(s) * \Delta \text{lk}_B(s); \\ & \Delta \text{lk}_A^+(s) * \Delta \text{lk}_B(s) \cup \Delta \text{lk}_A(s) * \Delta \text{lk}_B^+(s), \\ & \Delta \text{lk}_A^-(s) * \Delta \text{lk}_B(s) \cup \Delta \text{lk}_A(s) * \Delta \text{lk}_B^-(s)). \end{aligned}$$

However, for this to be correct, care has to be taken with regard to empty sets. We distinguish between the void complex $\emptyset$ with $X * \emptyset = \emptyset$ and the empty complex $\mathbb{S}^{-1}$ with $X * \mathbb{S}^{-1} = X$, which is consistent with $X * \mathbb{S}^{k-1}$ being the k -fold suspension of X . We set $\Delta \text{lk}_A^+(s) = \emptyset$, if and only if $A \cap p \cap r = \emptyset$. The justification for this is that we regard s as the tip of a cone over $\text{lk}_A(s)$ and want s to be in the cone over $\text{lk}_A^+(s)$ only if $A \cap p \cap r \neq \emptyset$. In the same way we set $\text{lk}_A^-(s) = \emptyset$ if and only if $A \cap p \subset q$.

With these conventions we obtain for $x \in p \cup (r \setminus q)$:

$$(4) \quad (\Delta \text{lk}_{\{x\}}(s); \Delta \text{lk}_{\{x\}}^+(s), \Delta \text{lk}_{\{x\}}^-(s)) \approx \begin{cases} (\mathbb{S}^0, \emptyset, \emptyset), & x \notin p, \\ (\mathbb{D}^0, \mathbb{S}^{-1}, \emptyset), & x \in p \cap q, \\ (\mathbb{D}^0, \emptyset, \mathbb{S}^{-1}), & x \in p \setminus r, \\ (\mathbb{D}^1, \{1\}, \{-1\}), & x \in p \cap (r \setminus q). \end{cases}$$

We take a look only at the most interesting case. For $x \in p \cap (r \setminus q)$ we have

$$(5) \quad \text{lk}_{\{x\}}(s) = \{(p^o, q \cup \{x\}, r^o), ((p \setminus \{x\})^o, q \cup \{x\}, r^o), \\ ((p \setminus \{x\})^o, q, r^o), ((p \setminus \{x\})^o, q, (r \setminus \{x\})^o), (p^o, q, (r \setminus \{x\})^o)\}$$

with two elements comparable if and only if they are written next to each other. Hence $\Delta \text{lk}_{\{x\}}(s)$ is an interval with boundary points $\text{lk}_{\{x\}}^+(s) = \{(p^o, q \cup \{x\}, r^o)\}$ and $\text{lk}_{\{x\}}^-(s) = \{(p^o, q, (r \setminus \{x\})^o)\}$.

From (4) one derives inductively using (3):

$$(6) \quad (\Delta \text{lk}_A(s); \Delta \text{lk}_A^+(s), \Delta \text{lk}_A^-(s)) \approx \begin{cases} (\mathbb{S}^{d_A}, \emptyset, \emptyset), & A \cap p = \emptyset, \\ (\mathbb{D}^{d_A}, \mathbb{S}^{d_A-1}, \emptyset), & \emptyset \neq A \cap p \subset q, \\ (\mathbb{D}^{d_A}, \emptyset, \mathbb{S}^{d_A-1}), & A \cap p \neq \emptyset, A \cap p \cap r = \emptyset, \\ (\mathbb{D}^{d_A}, \mathbb{D}_+^{d_A-1}, \mathbb{D}_-^{d_A-1}), & A \cap p \cap r \neq \emptyset, A \cap p \not\subset q \end{cases}$$

with $d_A = \#(A \cap (r \setminus q)) + \#(A \cap p) - 1$. Since $\text{lk}^+(s) = \text{lk}_{p \cup r}^+(s)$, $\text{lk}^-(s) = \text{lk}_{p \cup r}^-(s)$, $\text{lk}(s) = \text{lk}^+(s) \cup \text{lk}^-(s)$, the first case of (2) follows from the second case of (6), while the second case of (2) follows from the last case of (6). $\square$

4.11. Remark. The above Proposition will be used in the proof of Theorem 5.2 to show that $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ is the boundary of a regular neighbourhood of $\Delta\{\varphi \in F\text{Hom}(C_5 \setminus \{2,4\}, K_{n+2}) : \mathcal{C}\varphi(3) \subset \varphi(1)\}$. The proof of this special case would require exactly the same calculation, although the notation might possibly be more lucid. Indeed, the poset in (5) would become $\text{ind}(C_5 \setminus \{2,4\}) \setminus \{\emptyset\}$.

4.12. Proposition. *Let P be the face poset of a pl-triangulation of a compact manifold M^n . Let $B \subset P^o \times P \times P^o$ be given by*

$$B := \{(p^o, q, r^o) : q \leq r, p \cap r \neq \emptyset, p \not\leq q\}.$$

Then ΔB is homeomorphic to the boundary of a regular neighbourhood of the diagonal in $M \times M$.

Proof. We take up the notation of Proposition 4.10 and consider B as a subset of $P^o \times \text{Int } P$. We also define $D := \{(p^o, q, r^o) \in P^o \times \text{Int } P : p \leq q\}$. D is closed from above and hence ΔD is a full subcomplex of $\Delta(P^o \times \text{Int } P)$. $(\Delta N, \Delta B)$ is the simplicial neighbourhood of ΔD . By Proposition 4.10 ΔN is a manifold with boundary ΔB and hence a regular neighbourhood of ΔD by the Simplicial Neighbourhood Theorem 2.7. By Proposition 4.7 and Proposition 4.9 $\Delta D \subset \Delta(P^o \times \text{Int } P)$ is ambiently isotopic to the diagonal in $M \times M$. $\square$

5. $\text{Hom}(C_5, K_{n+2})$

We come back to our main object of study and collect the relevant results of the two preceding sections. First we can replace $\text{Hom}(C_5, K_{n+2})$ by a smaller complex.

5.1. Proposition. *Let $n \geq 0$ and consider $\{2,4\} \subset V(C_5)$. With the notation of Definition 3.1 we have $\text{Hom}(C_5, K_{n+2}) \approx \text{Hom}_{\{2,4\}}(C_5, K_{n+2})$.*

Proof. Since the independence complex of C_5 is also a 5-gon, this follows from Lemma 3.12. Alternatively it is easily checked directly that the precondition of Lemma 3.5 is satisfied, and we have done so in Example 3.8. $\square$

The graph $C_5 \setminus \{2, 4\}$ consists of an edge and an isolated vertex, and hence

$$\text{Hom}(C_5 \setminus \{2, 4\}, K_{n+2}) \approx \text{Hom}(K_2, K_{n+2}) \times \text{Hom}(K_1, K_{n+2}) \approx \mathbb{S}^n \times \mathbb{D}^{n+1}.$$

In Section 4 we have developed the techniques to describe the subcomplex $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$.

5.2. Theorem. *Let $n \geq 0$. Then $\text{Hom}(C_5, K_{n+2})$ is homeomorphic to the Stiefel manifold $\{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle = 0\}$.*

Proof. We examine $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$. The face poset of this complex is

$$\begin{aligned} \{\varphi : \{1, 3, 5\} \rightarrow \mathcal{P}(V(K_{n+2})) \setminus \{\emptyset\} : \\ \varphi(3) \cap \varphi(5) = \emptyset, \mathcal{C}(\varphi(1) \cup \varphi(3)) \neq \emptyset, \mathcal{C}(\varphi(5) \cup \varphi(3)) \neq \emptyset\}. \end{aligned}$$

The first condition comes from the edge $\{1, 5\}$, the second one ensures that φ can be extended to the vertex 2, the last one that it can be extended to 4. If we define P to be the poset $P := \mathcal{P}(V(K_{n+2})) \setminus \{\emptyset, V(K_{n+2})\}$ then it follows that the map

$$\begin{aligned} F\text{Hom}_{\{2,4\}}(C_5, K_{n+2}) &\rightarrow \{(p^o, q, r^o) \in P^o \times P \times P^o : q \leq r, p \not\leq q, p \cap r \neq \emptyset\} \\ \varphi &\mapsto (\mathcal{C}\varphi(3), \varphi(1), \mathcal{C}\varphi(5)) \end{aligned}$$

is an isomorphism. Since P is the face poset of the boundary of an $(n+1)$ -simplex, it follows from Proposition 4.12 that $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ is homeomorphic to the boundary of a regular neighbourhood of the diagonal in $\mathbb{S}^n \times \mathbb{S}^n$. Regular neighbourhoods are unique up to isotopy and $\{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle = 0\}$ is the boundary of the regular neighbourhood $\{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle \geq 0\}$ of the diagonal $\{(x, x) \in \mathbb{S}^n \times \mathbb{S}^n\}$. $\square$

Involutions. For the theorem of Babson and Kozlov mentioned in the introduction, the $\mathbb{Z}_2$ -action on $\text{Hom}(C_{2r+1}, K_{n+2})$ induced by an automorphism of C_{2r+1} that flips an edge is important. In [Cso05] the conjecture that $\text{Hom}(C_5, K_{n+2}) \approx \{(x, y) \in (\mathbb{S}^n)^2 : \langle x, y \rangle = 0\}$ is strengthened to a conjecture that there is a homeomorphism that takes the $\mathbb{Z}_2$ -action on $\text{Hom}(C_5, K_{n+2})$ to the action given by $(x, y) \mapsto (x, -y)$. In [Sch05] it was shown and used that there is an equivariant map between these spaces. This is essentially the map h that will appear in Theorem 5.4. We examine the constructions presented in this work more closely to show a homotopy version of the equivariant conjecture.

First we strengthen Proposition 4.12.

5.3. Proposition. *In the situation of Proposition 4.10, $(\Delta N, \Delta B)$ is a regular neighbourhood of the diagonal in $M \times M$.*

Proof. In Proposition 4.12 we have seen that ΔN is a regular neighbourhood of the subspace ΔD considered there and that there is an ambient isotopy moving ΔD to the diagonal. We will now show that this isotopy can be chosen to be the identity outside of $\Delta N \setminus \Delta B$.

The image of the isotopy constructed in the proof of Proposition 4.5 is contained in the union of the cells $\sigma \times \sigma$ for simplices σ of ΔP . We call this union the fat diagonal. Also every point in the image of the isotopy is contained in the interior

of one cell of the fat diagonal or in $\Delta(\text{Int } P)$. The complex $\Delta(P^\circ \times \text{Int } P)$ of Proposition 4.10 is a subdivision of $\Delta P \times \Delta P$ as discussed in Proposition 4.7. The subcomplex ΔN is contained in the fat diagonal, and no simplex of ΔB meets the interior of a cell of the fat diagonal. This proves the Proposition. $\square$

5.4. Theorem. *Let φ be an involution on $\text{Hom}(C_5, K_{n+2})$ induced by an automorphism of C_5 that flips an edge. Let $V := \{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle = 0\}$ and $\psi : V \rightarrow V$ be the involution $\varphi(x, y) = (x, -y)$. Then the following statements hold.*

- (i) *There is a homeomorphism $g : \text{Hom}(C_5, K_{n+2}) \xrightarrow{\cong} V$ with $\psi \circ g \simeq g \circ \varphi$.*
- (ii) *There is a homotopy equivalence $h : \text{Hom}(C_5, K_{n+2}) \xrightarrow{\cong} V$ with $\psi \circ h = h \circ \varphi$.*

Proof. Let us consider the automorphism of C_5 given by $j \mapsto 6 - j$. Figure 5 shows that the homeomorphism $\text{Hom}(C_5, K_{n+2}) \approx \text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ can be chosen to respect the induced involutions. In (7), $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ is identified with a subspace of $\Delta(P^\circ \times \text{Int } P)$ where P is the face poset of the boundary of the $(n+1)$ -simplex. This identification is compatible with the involution on $P^\circ \times \text{Int } P$ given by $(p^\circ, q, r^\circ) \mapsto (p^\circ, a(r), a(q)^\circ)$ where a is the map sending a face to the face opposite to it. On $\mathbb{S}^n \times \mathbb{S}^n \approx \Delta(P^\circ \times \text{Int } P)$ this map induces the identity on the first factor and the antipodal map on the second, and we will from now on consider the involution $(x, y) \mapsto (x, -y)$ on $\mathbb{S}^n \times \mathbb{S}^n$. By Proposition 5.3, $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ is the boundary of a regular neighbourhood of the diagonal in $\mathbb{S}^n \times \mathbb{S}^n$. The same is true of V , and for both spaces the involution on $\mathbb{S}^n \times \mathbb{S}^n$ exchanges the components of their complements. We define involutions

$$\begin{aligned} \Phi : [-1, 1] \times \text{Hom}(C_5, K_{n+2}) &\rightarrow [-1, 1] \times \text{Hom}(C_5, K_{n+2}), \\ (t, x) &\mapsto (-t, \varphi(x)), \end{aligned}$$

and

$$\begin{aligned} \Psi : [-1, 1] \times V &\rightarrow [-1, 1] \times V, \\ (t, x) &\mapsto (-t, \psi(x)). \end{aligned}$$

A structure of regular neighbourhood of the diagonal with boundary V defines for $\varepsilon > 0$ small enough an equivariant embedding of $[-1, 1] \times V$ in $\mathbb{S}^n \times \mathbb{S}^n$ which sends $\{-1\} \times V$ to the boundary of the ε -neighbourhood of the diagonal and $\{0\} \times V$ to V . Analogously and additionally using the homeomorphism $\text{Hom}(C_5, K_{n+2}) \approx \text{Hom}_{\{2,4\}}(C_5, K_{n+2})$, we obtain an embedding of $[-1, 1] \times \text{Hom}(C_5, K_{n+2})$ with the same image. This yields homeomorphisms

$$\begin{aligned} H : [-1, 1] \times \text{Hom}(C_5, K_{n+2}) &\xrightarrow{\cong} [-1, 1] \times V, \\ g : \text{Hom}(C_5, K_{n+2}) &\xrightarrow{\cong} V \end{aligned}$$

with

$$\begin{aligned} H(-1, x) &= (-1, g(x)) \quad \text{for all } x, \\ H \circ \Phi &= \Psi \circ H. \end{aligned}$$

It follows that

$$H(1, x) = \Psi(H(-1, \varphi(x))) = \Psi(-1, g(\varphi(x))) = (1, \psi(g(\varphi(x)))).$$

Therefore, if we denote by $p_V : [-1, 1] \times V \rightarrow V$ the projection, then $p_V \circ H$ is a homotopy between g and $\psi \circ g \circ \varphi$, which proves the first statement. If we define

$h(x) := p_V(H(0, x))$, then $h \simeq g$, since $g(x) = p_V(H(-1, x))$, and therefore h is a homotopy equivalence. Also

$$\begin{aligned} h(\varphi(x)) &= p_V(H(0, \varphi(x))) = p_V(H(\Phi(0, x))) = p_V(\Psi(H(0, x))) = \psi(p_V(H(0, x))) \\ &= \psi(h(x)), \end{aligned}$$

which proves the second statement. $\square$

5.5. Remark. The subspaces V and $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ of $\mathbb{S}^n \times \mathbb{S}^n$ are characteristic submanifolds of the involution on $\mathbb{S}^n \times \mathbb{S}^n$ in the language of [LdM71]. It might be possible to use the surgery techniques presented there to show that the induced involutions on them are equivalent, at least for $n \geq 3$ when the s -cobordism theorem is available. On the other hand the description of $\text{Hom}_{\{2,4\}}(C_5, K_{n+2})$ as a subpolyhedron of $\mathbb{S}^n \times \mathbb{S}^n$ that we have used is explicit enough to hope that it might be possible to prove this by a more direct approach.

REFERENCES

- [BK06a] BABSON, E. and KOZLOV, D. N. Complexes of graph homomorphisms. *Isr. J. Math.*, **152**:285–312, 2006. math.CO/0310056.
- [BK06b] —. Proof of the Lovász conjecture. *Annals of Mathematics*, 2006. In press, math.CO/0402395.
- [CL05] CSORBA, P. and LUTZ, F. H. Graph coloring manifolds, 2005. 22 pp., math.CO/0510177.
- [CLSW04] CSORBA, P., LANGE, C., SCHURR, I., and WASSMER, A. Box complexes, neighborhood complexes, and the chromatic number. *Journal of Combinatorial Theory, Series A*, **108**:159–168, 2004. math.CO/0310339.
- [Cso04] CSORBA, P. Homotopy types of box complexes, 2004. math.CO/0406118.
- [Cso05] —. *Non-tidy Spaces and Graph Colorings*. Ph.D. thesis, ETH Zürich, 2005.
- [Koz05a] KOZLOV, D. N. Cohomology of colorings of cycles, 2005. Preprint, math.AT/0507117.
- [Koz05b] —. A simple proof for folds on both sides in complexes of graph homomorphisms. *Proceedings of the AMS*, 2005. To appear, math.CO/0408262.
- [LdM71] LÓPEZ DE MEDRANO, S. *Involutions on Manifolds*, vol. 59 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1971.
- [Lov78] LOVÁSZ, L. Kneser's conjecture, chromatic number and homotopy. *J. Combinatorial Theory, Ser. A*, **25**:319–324, 1978.
- [MZ] MATOŮSEK, J. and ZIEGLER, G. M. Topological lower bounds for the chromatic number: A hierarchy. *Jahresbericht der DMV*. To appear, math.CO/0208072.
- [RS82] ROURKE, C. P. and SANDERSON, B. J. *Introduction to Piecewise-Linear Topology*. Springer-Verlag, 1982.
- [Sch05] SCHULTZ, C. A short proof of $w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = 0$ for all n and a graph colouring theorem by Babson and Kozlov, 2005. Preprint, 8pp., math.AT/0507346.
- [Sch06] —. Graph colourings, spaces of edges and spaces of circuits, 2006. Preprint, math.CO/0606763.
- [Seg68] SEGAL, G. Classifying spaces and spectral sequences. *Publ. Math. IHES*, **34**:105–112, 1968.
- [Wal88] WALKER, J. W. Canonical homeomorphisms of posets. *European J. Combin.*, **9**(2):97–107, 1988. ISSN 0195-6698.
- [Zee63] ZEEMAN, E. Unknotting of combinatorial balls. *Annals of Mathematics*, **78**:501–526, 1963.
- [Živ05] ŽIVALJEVIĆ, R. T. WI -posets, graph complexes and $\mathbb{Z}_2$ -equivalences. *Journal of Combinatorial Theory, Series A*, **111**:204–223, 2005. math.CO/0405419.

INSTITUT FÜR MATHEMATIK, MA 6-2, TECHNISCHE UNIVERSITÄT BERLIN, D-10623 BERLIN, GERMANY

E-mail address: carsten@codimi.de